

Anticancer evaluation of *Moringa oleifera* leaf extracts against lung cancer cell lines

Meenakshi Sharma¹, SB Sharma²

¹ Research Scholar, Faculty of Science, Motherhood University, Roorkee, Uttarakhand, India

² Supervisor, Professor and Dean, Faculty of Science, Motherhood University, Roorkee, Uttarakhand, India

Abstract

Moringa oleifera is a promising anticancer plants especially its leaf extracts having anticancer activities. In our study, we evaluate the anticancer activity of *Moringa oleifera* leaf extracts against A549 cell lines. The present investigate states that the percentage of cell inhibition of methanolic extract was found to be high against the A549 Cancer cell lines.

Keywords: *Moringa oleifera*, A549, cancer cell lines, methanolic extract

Introduction

There has been extensive interest in exploring the anticancer activities of various parts of the *M. oleifera* tree, and many published research articles describe promising results of in vitro and in vivo testing of various extracts from the Moringa plant. All parts of the *M. oleifera* tree have been tested for anticancer activity, including the leaves, seeds, bark, and roots. However, the most extensive research on the anticancer activities of *M. oleifera* has focused on the leaf extracts. The Moringa tree is an evergreen that grows new leaves year-round, with a projected production of six tons per hectare per year [1]. The leaves are rich in polyphenols and poly flavonoids, which are antioxidants and potential anticancer compounds [2]. Many researchers start by exploring the antioxidant activity and anti-inflammatory activity of the leaf extracts as a preliminary screening for anticancer activity. One of the factors that cause cancer is oxidative stress which is an imbalance in the production of free radicals and oxidants and their elimination by antioxidants [3]. Antioxidants can disrupt the formation of free radicals and reduce oxidative stress, which ultimately prevents cancer. The research showing the anticancer activities of moringa leaf extracts in vitro and the pathways/mechanisms of action through which these effects may occur.

Materials and methods

The *Moringa oleifera* leaf extracts from methanol, ethanol and acetone were used for testing the anticancer activity. The anticancer activity was performed against of cell lines namely A549 (Lung cancer cell line).

Anticancer activity of *Moringa oleifera* leaf extracts

Cell lines

The A549 (Lung cancer cell line) was obtained from National Centre for Cell Science (NCCS), Pune and grown in Eagles Minimum Essential Medium containing 10% fetal bovine serum (FBS). The cells were maintained at 37°C, 5% CO₂, 95% air and 100% relative humidity. Maintained cell cultures were passaged weekly, and the culture medium was changed twice a week.

Cell treatment procedure

The monolayer cells were detached with trypsin ethylene diamine tetraacetic acid (EDTA) to make single-cell suspensions. Viable cells were counted by trypan blue exclusion using a haemocytometer and diluted with medium containing 5% FBS to give the final density of 1x10⁵ cells/ml. One hundred microliters per well of cell suspension were seeded into 96-well plates at a plating density of 10,000 cells/well and incubated to allow for cell attachment at 37°C, 5% CO₂, 95% air and 100% relative humidity. After 24 h the cells were treated with serial concentrations of the test samples. They were initially dissolved in neat dimethyl sulfoxide (DMSO) and an aliquot of the sample solution was diluted to twice the desired final maximum test concentration with serum-free medium. Additional four serial dilutions were made to provide a total of five sample concentrations. Aliquots of 100 µl of these different sample dilutions were added to the appropriate wells already containing 100 µl of the medium, resulting in the required final sample concentrations. Following sample addition, the plates were incubated for an additional 48 h at 37°C, 5% CO₂, 95% air and 100% relative humidity. The medium containing without samples were served as control and triplicate were maintained for all concentrations.

MTT assay

MTT assay was performed based on the procedure described by (Mosmann, 1983). 3-[4, 5-dimethylthiazol-2-yl] 2, 5-diphenyltetrazolium bromide (MTT) is a yellow water-soluble tetrazolium salt. A mitochondrial enzyme in living cells, succinate- dehydrogenase, cleaves the tetrazolium ring, converting the MTT to an insoluble purple formazan. Therefore, the amount of formazan produced is directly proportional to the number of viable cells. After 48 h of incubation, 15µl of MTT (5mg/ml) in phosphate-buffered saline (PBS) was added to each well and incubated at 37°C for 4h. The medium with MTT was then discarded and the formed formazan crystals were solubilized in 100µl of DMSO and then measured the absorbance at 570 nm

using a microplate reader. The % cell inhibition was determined using the following formula.

$$\% \text{ Cell Inhibition} = 100 - \frac{\text{Abs (sample)}}{\text{Abs (control)}} \times 100.$$

Nonlinear regression graph was plotted between % Cell inhibition and Log concentration and IC₅₀ was determined using Graph Pad Prism software 8.

Results

Average percentage cell inhibition of Methanolic leaf extract

of *Moringa oleifera* leaf against A549 cells showed linear relationship, where the increased concentration of the extract increases the percentage of cell inhibition with an R2 value of about 0.968 (Figure 1). Ethanolic leaf extract of *Moringa oleifera* leaf against A549 cells showed a linear relationship, where the increased concentration of the extract increases the percentage of cell inhibition with an R2 value of about 0.975 (Figure 2). Acetone leaf extract of *Moringa oleifera* leaf against A549 cells showed a linear relationship, where the increased concentration of the extract increases the percentage of cell inhibition with an R2 value of about 0.973 (Figure.3)




Fig 1: Average percentage cell inhibition of methanolic leaf extract of *M. oleifera* against A549 cells




Fig 2: Average percentage cell inhibition of ethanolic leaf extract of *M. oleifera* against A549 cells

The 50% cell inhibition for the *Moringa oleifera* leaf extracts at different concentrations for the methanol, ethanol and acetone against A549 cells was determined. The cell inhibition was higher in the acetone leaf extract followed with the Methanolic leaf extract whereas it was observed to be very low in the ethanolic leaf extract (Figure. 2).

The percentage of cell inhibition at different concentrations like 50, 100, 150, 200, 250 and 300 $\mu\text{g/ml}$ against A549 cell line was calculated. At all concentrations, the percentage of cell inhibition was higher in the Methanolic extract with the range of about 17.94 ± 0.43 , 30.27 ± 1.16 , 48.07 ± 0.67 , 71.61 ± 0.76 , 82.32 ± 1.15 and 87.20 ± 1.00 with the p value of about <0.0007 at 50 $\mu\text{g/ml}$ and same p value of about <0.0001 for the other concentrations and different f values like 30.551, 54.199, 114.13, 167.66, 481.78 and 66.879. The cell inhibition was very low in the acetone extracts in all the concentrations with the range of

about 15.15 ± 0.35 , 18.16 ± 0.5 , 37.99 ± 0.67 , 55.03 ± 0.66 , 68.09 ± 0.73 and 75.85 ± 0.53 (Table 3). The inhibitory concentration (IC_{50}) for the leaf extract of *Moringa oleifera* against A549 cells in different solvents showed higher inhibitory concentration rate at acetone extract with the IC_{50} of about 193.73 ± 1.83 whereas it was very low in the ethanolic leaf extract with IC_{50} of about 175.21 ± 1.61 and it was found to be 178.78 ± 1.2 in the Methanolic leaf extract with the p-value of about <0.0001 and f value of about 161.22 (Table.4). The significance rate of percentage of cell inhibition at different concentrations for the different solvent extracts of *Moringa oleifera* leaf against A549 cells was determined. At the concentration of 50 $\mu\text{g/ml}$, it was highly significant with methanol and acetone whereas non-significant in ethanol with acetone. 100, 150 and 200 $\mu\text{g/ml}$ showed highly significant with all the solvents like methanol with ethanol and acetone, ethanol with acetone (Table. 5).




Fig 3: Average percentage cell inhibition of acetone leaf extract of *M. oleifera* against A549 cells




Fig 4: Fifty percent inhibition (IC_{50}) concentration of *M. oleifera* crude extract with different solvents for A549 cells

Table 1: Percentage cell inhibition of different solvent leaf extract of *M. oleifera* ($\mu\text{g/ml}$) against A549 Cells

Concentrations	Methanol	Ethanol	Acetone	p value	f value
50	17.94 \pm 0.43	16.02 \pm 0.25	15.15 \pm 0.35	<0.0007	30.551
100	30.27 \pm 1.16	22.13 \pm 0.64	18.16 \pm 0.51	<0.0001	54.199
150	48.07 \pm 0.67	41.96 \pm 0.73	37.99 \pm 0.67	<0.0001	114.13
200	71.61 \pm 0.76	61.81 \pm 0.35	55.03 \pm 0.66	<0.0001	167.66
250	82.32 \pm 1.15	75.98 \pm 0.43	68.09 \pm 0.73	<0.0001	481.78
300	87.20 \pm 1.00	82.39 \pm 1.12	75.85 \pm 0.53	<0.0001	66.879
p value	< 0.0001	< 0.0001	< 0.0001		
f value	379.71	671.53	569.88		

Table 2: Fifty percent inhibition (IC_{50}) concentration of *M. oleifera* crude extract with different solvents for A549 cells

Solvents	IC_{50} ($\mu\text{g/ml}$)
Methanol	162.61 \pm 13.15
Ethanol	175.21 \pm 1.61
Acetone	193.73 \pm 1.83
p value	< 0.0001
f value	161.22

Table 3: Comparative significance percentage cell inhibition of different solvent leaf extract of *M. oleifera* ($\mu\text{g/ml}$) against A549 Cells

Solvents	Ethanol	Acetone	Ethanol	Acetone	Ethanol	Acetone
	50		100		150	
Methanol	**	***	**	***	***	***
Ethanol		ns		*		**
	200		250		300	
Methanol	**	***	**	***	**	***
Ethanol		***		***		**

Discussion

Though *Moringa oleifera* was established for its prospective medicinal values, more researches have to be carried out to study its anticancer activities. Hence, the methanol, ethanol and acetone extracts of *Moringa oleifera* leaf was evaluated for its cytotoxicity activities. The presence of many organosulphur compounds in the *Moringa oleifera* leaves was highly responsible for the prevention of the cancer cells (Aja, *et al.*, 2014). Hence as per the earlier report, the present investigations revealed that the cell inhibition increases by increased concentrations of the *Moringa oleifera* leaf extract with different solvents (Figure 1- 4). The *in vivo* study of the methanolic and ethanolic extracts of *Moringa oleifera* leaf delays the cancer cell proliferation in various cancer cells and increases the survival rate of the cancer patients (Purwal, *et al.*, 2010). Hence the present investigation also states that the percentage of cell inhibition of methanolic extract was found to be high against the A549 cell lines. Recent researches suggested that the compounds like niazimicin and the isothiocyanate present in the ethanolic extract of the *Moringa oleifera* leaf possessed to contain the antioxidant and anticancer activities against many cancer cell lines like breast, lung and colorectal cancers (Hermawan, *et al.*, 2012).

Acknowledgement: Authors wants to thanks for help and support.

Conflict of Interest: Authors shows no conflict of interest

References

1. Abdulaziz Rabi Abdulkadir, Dhiya Dalila Zawawi, Sarwar Jahan MD. DPPH antioxidant activity, total phenolic and total flavonoid content of different part of Drumstic tree (*Moringa oleifera* Lam). J. Chem. Pharm. Res. 2015; 7(4):1423-1428
2. Abdulrahman KA, Sulaiman MA, Tanwir Athar MD, Abdul QK, Hamoud S. *Moringa oleifera* as an Anti-Cancer Agent against Breast and Colorectal Cancer Cell Lines. Journal. Pune. 01, 2015.
3. Adam Hermawan, Kholid Alfian Nur, Sarmoko Dyaningtyas Dewi, Pamungkas Putri, Edy Meiyanto. Ethanolic extract of *Moringa oleifera* increased cytotoxic effect of doxorubicin on HeLa cancer cells. Journal of Natural Remedies. 2012; 12(2):108-114.
4. Aja PM, Nwachukwu N, Ibiam UA, Igwenyi IO, Ofor CE, Orji UO *et al.* Chemical Constituents of *Moringa oleifera* Leaves and Seeds from Abakaliki, Nigeria. American Journal of Phytomedicine and clinical therapeutics. 2014; 2(3):310-321.
5. Amaglo NK, Bennet RN, Lo Curto RB, Rosa EAS, Crea F. Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* Lam. grown in Ghana. Food. Chem. 2010; 122:1047-1054.
6. Hermawan A, Nur KA, Sarmoko D, Dewi P, Putri E, Meiyanto. Ethanolic extract of *Moringa oleifera* increased cytotoxic effect of doxorubicin on HeLa cancer cells. J. Nat. Remedies. 2012; 12:108-114